

Publications

- Velegraki, M., **Papakonstanti, E.A.**, Mavroudi, I., Psyllaki, M., Tsatsanis, C., and Papadaki, E.A. Impaired clearance of apoptotic cells leads to HMGB1 release in the bone-marrow of MDS patients and induces TLR4-mediated cytokine production. (2012), (*Hematologica*, in revision).
- Tzenaki, N., Vergetaki, A., Zwaenepoel O., **Papakonstanti E.A.** FAK links RhoA with PTEN regulating migration and cell survival (2012, submitted).
- Vergetaki, A., Jeschke, U., Vrekoussis, T., Sabatini, L., Taliouri, E., Christoforaki, V., **Papakonstanti, E.A.**, Makrigiannakis, A. Overexpression of CRH, UCN, CRHR1 AND CRHR2 in ectopic endometrium of women with endometriosis (*J Clin Endocrinol Metab*, under review).
- Tzenaki, N., Andreou, M., Stratigi, K., Vergetaki, A., Makrigiannakis, A., Vanhaesebroeck, B., **Papakonstanti, E.A.** High levels of p110 δ PI3K expression in solid tumour cancer cells suppresses PTEN activity, generating cellular sensitivity to p110 δ inhibitors through PTEN activation. *FASEB J* (2012) 26(6), 2498-2508.
- Zwaenepoel O., Tzenaki, N., Vergetaki, A., Makrigiannakis, A., Vanhaesebroeck B., **Papakonstanti E.A.** Functional CSF-1 receptors are located at the nuclear envelope and activated via the p110 δ isoform of PI 3-kinase. *FASEB J* (2012) 26 (2), 691-706.
- Papadopoulou N., **Papakonstanti E.A.**, Kallergi G., Alevizopoulos K., Stournaras C. [Membrane androgen receptor activation in prostate and breast tumor cells: molecular signaling and clinical impact.](#) *IUBMB Life* (2009) 61, 56-.
- **Papakonstanti E.A.**, Zwaenepoel O., Bilancio A., Burns E., Houseman B., Shokat K., Ridley A.J., Vanhaesebroeck B. Distinct roles of class IA PI 3-kinase isoforms in primary and immortalised macrophages. *J Cell Science* (2008) 121, 4124-33.
- **Papakonstanti E.A.**, and Stournaras C. Cell responses regulated by early reorganization of actin cytoskeleton. *FEBS Lett.* (2008) 582, 2120-7.
- **Papakonstanti E.A.**, Ridley A.J., Vanhaesebroeck B. The p110 δ isoform of PI 3-kinase negatively controls RhoA and PTEN. *EMBO J* (2007) 26, 3050-61.
- Eickholt B.J., Ahmed A.I., Davies M., **Papakonstanti E.A.**, Pearce W., Starkley M.L., Bilancio A., Need A.C., Smith A.J.H., Hall S.M., Hamers F.P., Giese K.P., Bradbury E.J., Vanhaesebroeck B. Control of axonal growth and regeneration of sensory neurons by p110 δ PI 3-kinase. *PLoS ONE* (2007) 2, e869.
- **Papakonstanti E.A.**, and Stournaras C. Actin Cytoskeleton Architecture and Signaling in Osmosensing. *Methods Enzymol* (2007) 428, 227-40.
- Kampa M, Kogia C, Theodoropoulos P A, Anezinis P, Charalampopoulos I, **Papakonstanti E A**, Stathopoulos E N, Hatzoglou A, Stournaras C, Gravanis A, Castanas E. Activation of membrane androgen receptors potentiates the antiproliferative effects of paclitaxel on human prostate cancer cells. *Mol. Cancer Therapeutics* (2006) 5, 1342-51.
- Hatzoglou A., Kampa M., Kogia C., Charalampopoulos I., Theodoropoulos P.A., Anezinis P., Dambaki C., **Papakonstanti E.A.**, Stathopoulos E.N., Stournaras C., Gravanis A. and Castanas E. Membrane androgen receptor activation induces apoptotic regression of human prostate cancer cells *in vitro* and *in vivo*. *J Clin. Endocrinol Metab.* (2005) 90, 893-903.
- **Papakonstanti E.A.**, and Stournaras C. TNF α promotes survival of OK cells via Cdc42-induced PLC- γ 1 activation and actin filament redistribution. *Mol Biol Cell* (2004) 15, 1273-86.
- Kampa M., **Papakonstanti E.A.**, Alexaki V-I., Hatzoglou A., Stournaras C. and Castanas E. The opioid agonist ethylketocyclazocine reverts the rapid, non-genomic, effects of membrane testosterone receptors in the human prostate LNCaP cell line. *Exp. Cell Research* (2004) 294, 434-45.
- **Papakonstanti E.A.**, Kampa M., Castanas E., and Stournaras C. A rapid, non-genomic, signaling pathway regulates the actin reorganization induced by membrane testosterone receptors' activation. *Mol. Endo* (2003) 17, 870-81.

- Kallergi G., Tsapara A., Kampa M., **Papakonstanti E.A.**, Krasagakis K., Castanas E., and Stournaras C. Distinct signaling pathways regulate differential opioid effects on actin cytoskeleton in malignant MCF7 and nonmalignant MCF12A human breast epithelial cells. *Exp. Cell Res.* (2003) 288, 94-109.
- Kampa M., **Papakonstanti E.A.**, Hatzoglou A., Stathopoulos S., Stournaras C. and Castanas E. The human prostate cancer cell line LNCaP bears functional membrane testosterone receptors, which increase PSA secretion and modify actin cytoskeleton. *FASEB J* (2002) 16, 1429-1431.
- **Papakonstanti E.A.**, and Stournaras C. Association of PI-3 kinase with PAK1 leads to actin phosphorylation and cytoskeletal reorganization. *Mol Biol Cell* (2002) 13, 2946-2962.
- **Papakonstanti E.A.**, Emmanouel D.S., Gravanis A., and Stournaras C. PLC- γ 1 signalling pathway and villin activation are involved in actin cytoskeleton reorganization induced by the Na⁺/P_i cotransport upregulation. *Molecular Medicine* (2000) 6, 303-318.
- **Papakonstanti E.A.**, Vardaki E.A., and Stournaras C. Actin cytoskeleton: a signalling sensor in cell volume regulation. *Cell. Physiol. Biochem.* (2000) 10, 257-264.
- Koukouritaki S.B., Vardaki E.A., **Papakonstanti E.A.**, Lianos E., Stournaras C., and Emmanouel D.S. TNF- α induces actin cytoskeleton reorganization in glomerular epithelial cells involving tyrosine phosphorylation of paxillin and focal adhesion kinase. *Molecular Medicine* (1999) 5, 383-393.
- Panagiotou, S., Bakogeorgou, E., **Papakonstanti, E.A.**, Hatzoglou, A., Wallet, F., Dussert, C., Stournaras, C., Martin, P.M. and Castanas, E. Opioid agonists modify breast cancer cell proliferation by blocking cells to the G₂/M phase of the cycle: Involvement of cytoskeletal elements. *J. Cell. Biochem.* (1999) 73, 204-211.
- **Papakonstanti, E.A.**, Bakogeorgou, E., Castanas, E., Emmanouel, D.S., Hartig, R. and Stournaras, C. Early alterations of actin cytoskeleton in OK cells by opioids. *J. Cell. Biochem.* (1998) 70, 60-69.
- Hatzoglou, A., Bakogeorgou, E., **Papakonstanti, E.A.**, Stournaras, C., Emmanouel, D.S. and Castanas, E. Identification and characterization of opioid and somatostatin binding sites in the Opossum Kidney (OK) cell line, and their effect on growth. *J. Cell. Biochem.* (1996) 63, 410-421.
- **Papakonstanti, E.A.**, Emmanouel, D.S., Gravanis, A. and Stournaras, C. Na⁺/P_i co-transport alters rapidly cytoskeletal protein polymerization dynamics in opossum kidney cells. *Biochem. J.*, (1996) 315, 241-247.